

A BOUND FOR THE PERIMETER OF INNER PARALLEL BODIES

SIMON LARSON

ABSTRACT. We provide a sharp lower bound for the perimeter of the inner parallel sets of a convex body Ω . The bound depends only on the perimeter and inradius of the original body and states that

$$|\partial\Omega_t| \geq \left(1 - \frac{t}{R_{in}}\right)_+^{n-1} |\partial\Omega|.$$

In particular the bound is independent of any regularity properties of $\partial\Omega$. As a by-product of the proof we establish precise conditions for equality. The proof, which is rather straightforward, is based on the construction of an extremal set for a certain optimization problem and the use of basic properties of mixed volumes.

1. INTRODUCTION

Given a convex domain $\Omega \subset \mathbb{R}^n$ we consider the family of its *inner parallel sets*. We denote by Ω_t the inner parallel set at distance $t \geq 0$, which is defined by

$$\Omega_t = \{x \in \mathbb{R}^n \mid \text{dist}(x, \Omega^c) \geq t\} = \Omega \sim tB.$$

Here B is the unit ball in \mathbb{R}^n and \sim denotes the *Minkowski difference*; a precise definition is given in section 1.1. Correspondingly, the *outer parallel set* at distance $t \geq 0$ is the set

$$\{x \in \mathbb{R}^n \mid \text{dist}(x, \Omega) \leq t\} = \Omega + tB,$$

where $+$ denotes the *Minkowski sum*. The main goal of this paper is to provide a lower bound on the perimeter of Ω_t in terms of the perimeter of Ω .

The theory of parallel bodies had one of its early breakthroughs in Steiner's proof that the volume of an outer parallel body $\Omega + tB$ can be expressed as a polynomial in t . This statement is the so-called *Steiner formula*

$$|\Omega + tB| = \sum_{i=0}^n \binom{n}{i} t^i W_i(\Omega).$$

The coefficients W_i of the polynomial are the *quermassintegrals* of Ω . The set of quermassintegrals contains several important geometric quantities; for instance we have that $W_0(\Omega) = |\Omega|$ and $nW_1(\Omega) = |\partial\Omega|$.

The Steiner formula inspired the development of the Brunn-Minkowski theory which is a central part of classic convex geometry. This theory arises when one combines the concepts of Euclidean volume and Minkowski addition. A key concept in the study of Brunn-Minkowski theory, which will be of importance in what follows, is that of *mixed volumes*. The mixed volume W is defined as the n -linear functional over \mathcal{K}_0^n (see definition in section 1.1) which for

Date: 20th January, 2019.

2010 Mathematics Subject Classification. Primary 52A20, 52A38; Secondary 52A40, 52A39.

Key words and phrases. Convex geometry, perimeter, inner parallel sets, mixed volumes, form body, inradius.

any $\Omega_1, \dots, \Omega_k \in \mathcal{K}_0^n$ and all $\lambda_1, \dots, \lambda_k \geq 0$ satisfies

$$|\lambda_1 \Omega_1 + \dots + \lambda_k \Omega_k| = \sum_{i_1=1}^k \dots \sum_{i_n=1}^k \lambda_{i_1} \dots \lambda_{i_n} W(\Omega_{i_1}, \dots, \Omega_{i_n}).$$

From this characterization one can show that the mixed volume of an n -tuple of non-empty convex bodies is uniquely determined and that W is well-defined as a functional from $(\mathcal{K}_0^n)^n$ to \mathbb{R} , see for instance [16].

The mixed volumes generalize the quermassintegrals that appear as coefficients in the Steiner polynomial above. Comparing the Steiner formula and the above identity one finds that

$$W_i(\Omega) = W(\underbrace{\Omega, \dots, \Omega}_{n-i}, \underbrace{B, \dots, B}_i).$$

We will in section 1.1 present some elementary properties of the mixed volumes that will be needed in the sequel. For further details and a comprehensive overview of the Brunn-Minkowski theory and mixed volumes we refer to [16] and references therein.

Expressions similar to that of the Steiner formula can be obtained for the mixed volumes of Ω and B , which, as mentioned above, are simply the quermassintegrals of Ω . The resulting identities are called the *Steiner formulae*:

$$W(\underbrace{\Omega, \dots, \Omega}_{n-k}, \underbrace{B, \dots, B}_k) = \sum_{i=0}^{n-k} \binom{n-k}{i} W(\underbrace{\Omega, \dots, \Omega}_{n-k-i}, \underbrace{B, \dots, B}_{k+i}).$$

With this in hand the study of outer parallel sets reduces to applying the extensive theory of Minkowski addition and mixed volumes.

The Steiner formulae and the surrounding theory has influenced several areas of modern mathematics. Important applications may be found in the work of Federer on curvature measures in geometric measure theory (see [7]) and Weyl's tube formula in differential geometry (see [21]). The Steiner formula also appears in many other fields, for instance algebraic and stochastic geometry, geometric functional analysis and of course both convex geometry and analysis (see for instance [5, 11, 18]).

The concept of Minkowski differences may in some sense be viewed as the inverse operation to addition of convex sets, but it is far from being as well behaved as the Minkowski sum. One of the main obstacles in the development of a similar theory as for the addition is that the Minkowski difference is (in contrast to the Minkowski sum) not a vectorial operation. Moreover, the regularity properties of $\partial\Omega_t$ may be very different from those of $\partial\Omega$. Both of these properties are demonstrated, for the two-dimensional case, in Figure 1. Nonetheless, and one may argue that it is due to this increased complexity, the theory of inner parallel sets is rich and has many beautiful connections and applications in both convex geometry and analysis (see for instance [2, 4, 12, 13, 14, 15]).

For the Minkowski difference there exists, in general, no counterpart to the Steiner formula. It was conjectured by Matheron in [13] that the volume of a Minkowski difference can be controlled using the *alternating Steiner polynomial*. That is, for Ω and $\tilde{\Omega}$ convex bodies in \mathbb{R}^n he conjectured that

$$|\Omega \sim t\tilde{\Omega}| \geq \sum_{i=0}^n \binom{n}{i} (-t)^i W(\underbrace{\Omega, \dots, \Omega}_{n-i}, \underbrace{\tilde{\Omega}, \dots, \tilde{\Omega}}_i).$$

However this turns out to be false. The conjecture was disproved by Hernández Cifre and Saorín in [9]. Further, the same authors obtain in [10] bounds on the quermassintegrals of inner parallel sets in a more general setting than that described above. The setting considered in [10] is called




FIGURE 1. The inner parallel set of some convex bodies in \mathbb{R}^2 . Note that $\tilde{\Omega}_t$ is equal to $\hat{\Omega}_t$ even though $\tilde{\Omega} \neq \hat{\Omega}$.

the *relative parallel bodies* of Ω (see for instance [2, 6, 14]). The relative parallel bodies are obtained by, instead of adding or subtracting (in Minkowski sense) tB , one adds or subtracts tE for some set $E \in \mathcal{K}_0^n$. The family of sets obtained are called the parallel bodies of Ω relative to E . The inequalities obtained in this article are closely related to those in [10] and using similar techniques the results here could, at least in some sense, be generalized to relative parallel bodies. However, in these generalizations one loses the connection to the spectral theoretic applications that motivated the work in this paper.

One of the classical results obtained from the Steiner formula is the relation between the perimeter of Ω and that of its outer parallel sets. Using this result one can obtain the following theorem, see [19]. The main result of this paper is an improvement of statement (ii) of the theorem.

Theorem 1.1 (Modified Steiner inequality [19]). *Let Ω be a convex domain in \mathbb{R}^n with volume $|\Omega|$ and surface area $|\partial\Omega|$ such that the curvature of $\partial\Omega$ is at each point bounded from above by $1/K$ for some $K > 0$. Then the following bounds hold for the inner parallel sets of Ω*

$$(i) \quad |\Omega_t| = |\Omega| - \int_0^t |\partial\Omega_\tau| d\tau \text{ for all } 0 \leq t \leq K,$$

$$(ii) \quad |\partial\Omega_t| \geq |\partial\Omega| \left(1 - \frac{(n-1)}{K}t\right),$$

$$(iii) \quad |\Omega_r| \geq |\Omega_s| \text{ for all } r \leq s,$$

$$(iv) \quad |\partial\Omega_r| \geq |\partial\Omega_s| \text{ for all } r \leq s.$$

Further, the curvature at each point of $\partial\Omega_t$ is bounded from above by $(K-t)^{-1}$ for any $0 \leq t < K$.

Our study of this problem is motivated by the work of Geisinger, Laptev and Weidl in [8]. In one part of this article the authors use statement (ii) of the above theorem to obtain improved bounds on Riesz means of eigenvalues for the Dirichlet Laplacian on a convex domain in \mathbb{R}^n . Furthermore, for convex domains in the plane that satisfy the inequality

$$(1) \quad |\partial\Omega_t| \geq \left(1 - \frac{3t}{\omega}\right)_+ |\partial\Omega|,$$

where ω denotes the width of Ω , the authors improve further upon these bounds. However these particular results of [8] are conditional on that (1) holds. The authors conjecture that this is the case for any planar convex body but do not provide a proof. In this paper we prove that the bound holds for any convex set in \mathbb{R}^2 and that similar bounds hold in arbitrary dimension.

The main result of this paper is contained in the next theorem. In an upcoming paper the techniques of [8] will be combined with this result to obtain further geometrical improvements of Berezin type bounds for the Dirichlet eigenvalues of the Laplacian on convex domains.

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^n$ be a convex domain with inner radius R_{in} . Then, for any inner parallel set Ω_t , $t \geq 0$, it holds that*

$$|\partial\Omega_t| \geq \left(1 - \frac{t}{R_{in}}\right)_+^{n-1} |\partial\Omega|.$$

Further, equality holds for some $t \in (0, R_{in})$ if and only if Ω is homothetic to its form body. If this is the case equality holds for all $t \geq 0$.

Using the above theorem and some known bounds for the inner radius and width of convex bodies we are able to conclude that the conjectured inequality (1) holds and provide the following generalization to higher dimensions.

Corollary 1.3. *Let $\Omega \subset \mathbb{R}^n$ be a convex domain with width ω . Then, for the inner parallel sets of Ω we have that*

$$\begin{aligned} |\partial\Omega_t| &\geq \left(1 - \frac{2\sqrt{n}}{\omega} t\right)_+^{n-1} |\partial\Omega| && \text{if } n \text{ is odd} \\ |\partial\Omega_t| &\geq \left(1 - \frac{2(n+1)}{\omega\sqrt{n+2}} t\right)_+^{n-1} |\partial\Omega| && \text{if } n \text{ is even.} \end{aligned}$$

In both cases equality holds if Ω is a regular $(n+1)$ -simplex.

We conjecture that equality holds if and only if Ω is the regular simplex but are not currently able to provide a proof. That this holds in \mathbb{R}^2 is clear but the generalization of the proof to higher dimensions eludes us.

The result developed here is in several aspects an improvement of the corresponding statement in Theorem 1.1. Firstly, the assumptions on Ω are substantially less restrictive in our statement. We have no other conditions than convexity whilst the earlier result requires boundedness of the curvature of $\partial\Omega$. Further, by noting that

$$\left(1 - \frac{t}{K}\right)_+^{n-1} \geq \left(1 - \frac{(n-1)}{K} t\right)_+$$

and that the maximum curvature of the boundary of a convex set is always larger than the reciprocal of the inradius of the set one can conclude that Theorem 1.2 implies the bound from Theorem 1.1. We note that if the distance t is less than the maximal curvature of $\partial\Omega$ the problem is somewhat less complex, since if this is the case the outer parallel set of Ω_t at distance t is precisely the set Ω (which is not the case in general).

1.1. Notation and preliminaries. Let \mathcal{K}_0^n denote the set of all convex bodies in \mathbb{R}^n that have nonempty interior. Throughout the paper Ω will belong to \mathcal{K}_0^n . Also let B denote the closed unit ball in \mathbb{R}^n and let \mathbb{S}^{n-1} denote the corresponding sphere. A closed ball of radius r centred at $x \in \mathbb{R}^n$ is denoted by $B_r(x)$. For notational simplicity we denote both volume and surface measure by $|\cdot|$. This will appear in two forms, the volume of a set $|\Omega|$ and the surface measure of its boundary $|\partial\Omega|$. Further, we will make use of the notation $x_{\pm} = (|x| \pm x)/2$.

For two sets $K, L \in \mathcal{K}_0^n$ the *Minkowski sum* $(+)$ and *difference* (\sim) are defined by

$$\begin{aligned} K + L &:= \{x + y : x \in K, y \in L\}, \\ K \sim L &:= \{x \in \mathbb{R}^n : x + L \subseteq K\}. \end{aligned}$$

It is a direct consequence of the definitions that we, as claimed in the introduction, equivalently can define the inner parallel body Ω_t , $t \geq 0$, as $\Omega \sim tB$ [16]. Similarly the outer parallel body can be written as $\Omega + tB$.

The *inradius* R_{in} of a set $\Omega \in \mathcal{K}_0^n$ is defined as the radius of the largest ball contained in Ω , or equivalently (see for instance [16]) as

$$R_{in} = \sup\{r > 0 : \Omega \sim rB \neq \emptyset\}.$$

The observation contained in the next lemma is intuitively clear but of central importance in what follows. The main consequence of the lemma is the fact that the inradius of Ω_t behaves very nicely with respect to t .

Lemma 1.4. *Let $\Omega \in \mathcal{K}_0^n$ have inradius R_{in}^0 . Then, for any $t \in [0, R_{in}^0]$ the inradius R_{in}^t of Ω_t satisfies*

$$R_{in}^t = R_{in}^0 - t.$$

Proof. Let $x_0 \in \mathbb{R}^n$ be such that $R_{in}^0 B(x_0) \subseteq \Omega$. For each $x \in R_{in}^0 B(x_0)$ we have that $\text{dist}(x, \partial\Omega) \geq \text{dist}(x, \partial R_{in}^0 B(x_0))$ and hence it holds that

$$\Omega_t \supseteq (R_{in}^0 B(x_0))_t = (R_{in}^0 - t)B(x_0).$$

We conclude that the inequality $R_{in}^t \geq R_{in}^0 - t$ holds. To prove the reverse inequality we observe that for $\tilde{x}_0 \in \mathbb{R}^n$ such that $R_{in}^t B(\tilde{x}_0) \supseteq \Omega_t$ we have, by the definition of Ω_t , that $\text{dist}(R_{in}^t B(\tilde{x}_0), \partial\Omega) \geq t$. Hence it holds that

$$(R_{in}^t + t)B(\tilde{x}_0) = R_{in}^t B(\tilde{x}_0) + tB \supseteq \Omega,$$

which implies the inequality $R_{in}^0 \geq R_{in}^t + t$. The statement of the lemma follows. \square

A classical result in convex geometry is that the volume of a Minkowski sum $\lambda_1 K_1 + \dots + \lambda_m K_m$ is, for $\lambda_1, \dots, \lambda_m \geq 0$ and $K_1, \dots, K_m \in \mathcal{K}_0^n$, a homogeneous n -th degree polynomial in the λ_i (see [3, 16, 20]). That is we can write

$$|\lambda_1 K_1 + \dots + \lambda_m K_m| = \sum_{i_1=1}^m \dots \sum_{i_n=1}^m \lambda_{i_1} \dots \lambda_{i_n} W(K_{i_1}, \dots, K_{i_n}).$$

The coefficients $W(K_{i_1}, \dots, K_{i_n})$ are the mixed volumes of K_1, \dots, K_m . In what follows we will use several properties of W . We list the properties here, and for proofs we refer to [3, 16]:

- The mixed volume is a symmetric functional on $(\mathcal{K}_0^n)^n$. For $K_1, \dots, K_n \in \mathcal{K}_0^n$ and $\sigma \in S_n$ it holds that

$$W(K_1, \dots, K_n) = W(K_{\sigma(1)}, \dots, K_{\sigma(n)}).$$

- W is Minkowski linear in each of its arguments:

$$W(\lambda K + \lambda' K', K_2, \dots, K_n) = \lambda W(K, K_2, \dots, K_n) + \lambda' W(K', K_2, \dots, K_n).$$

- W is monotonically increasing with respect to inclusions: if $K' \subseteq K$ then

$$W(K', K_2, \dots, K_n) \leq W(K, K_2, \dots, K_n).$$

- The mixed volume is invariant under translations in each argument. For $K_1, \dots, K_n \in \mathcal{K}_0^n$ and any $x_1, \dots, x_n \in \mathbb{R}^n$ we have that

$$W(K_1, \dots, K_n) = W(x_1 + K_1, \dots, x_n + K_n).$$

- The perimeter of $K \in \mathcal{K}_0^n$ is, up to a constant, equal to a mixed volume:

$$|\partial K| = nW(B, K, \dots, K).$$

We will by $h(K, u)$ denote the *support function* of $K \in \mathcal{K}_0^n$ which is defined for any $u \in \mathbb{R}^n$ as

$$h(K, u) = \sup\{\langle x, u \rangle : x \in K\}.$$

The restriction of $h(K, u)$ to $u \in \mathbb{S}^{n-1}$ reduces to the function describing the distance from the origin to the supporting plane of K with normal u . The existence of a unique such plane is clear from convexity. We will in what follows denote such a supporting plane by $H(K, u)$. In this notation we have the following characterization of the supporting planes of Ω :

$$H(K, u) = \{x \in \mathbb{R}^n : \langle x, u \rangle = h(K, u)\}.$$

The following properties of $h(K, u)$ will be needed in the sequel:

- For any $K, L \in \mathcal{K}_0^n$ and $\alpha, \beta > 0$ it holds that

$$h(\alpha K + \beta L, u) = \alpha h(K, u) + \beta h(L, u).$$

- For any $u \in \mathbb{S}^{n-1}$ and $K, L \in \mathcal{K}_0^n$ it holds that

$$h(K \sim L, u) \leq h(K, u) - h(L, u).$$

- For $x \in \partial(K \sim L)$ there exists a normal vector u of $\partial(K \sim L)$ at x such that

$$h(K \sim L, u) = h(K, u) - h(L, u).$$

Proofs of the above properties and further results connected to the support function can be found in most literature on convex geometry, see for instance [3, 16, 20].

The *width* ω of $\Omega \in \mathcal{K}_0^n$ is defined as

$$\omega = \inf\{h(\Omega, u) + h(\Omega, -u) : u \in \mathbb{S}^{n-1}\}.$$




FIGURE 2. The width and inradius of a convex body in \mathbb{R}^2 .

A point $x \in \partial\Omega$ is called *regular* if the supporting plane at x is uniquely defined, that is if there is a unique $u \in \mathbb{S}^{n-1}$ such that

$$x \in H(\Omega, u) \cap \Omega.$$

The set of all regular points of $\partial\Omega$ is denoted by $\text{reg}(\Omega)$. Further, we call $u \in \mathbb{S}^{n-1}$ an *extreme normal* of Ω if u is the unique normal of $\partial\Omega$ at some $x \in \text{reg}(\Omega)$ and let $\mathcal{U}_0(\Omega)$ denote the set of extreme normals of Ω .

In the proof of Theorem 1.2 we will need the concept of the *form body* Ω^* of $\Omega \in \mathcal{K}_0^n$, which is (see [10]) defined as

$$\Omega^* = \bigcap_{u \in \mathcal{U}_0(\Omega)} \{x \in \mathbb{R}^n : \langle x, u \rangle \leq 1\}.$$

If Ω is a polytope then Ω^* is the polytope that has the same set of normals as Ω , but with each face translated so that it is tangent to the unit ball. If instead the boundary of Ω is smooth (every point is regular) then $\Omega^* = B$.




FIGURE 3. The form body of two convex sets in \mathbb{R}^2 .

The following lemma will be needed in our main proof and is an almost direct consequence of the definitions of Ω^* and R_{in} combined with the fact that almost every point of $\partial\Omega$ is regular.

Lemma 1.5. *Let $\Omega \in \mathcal{K}_0^n$ have inradius R_{in} . Then there exists $x \in \mathbb{R}^n$ such that $x + R_{in}\Omega^* \subseteq \Omega$.*

2. PROOF OF THE MAIN RESULT

The idea of the proof is to construct, for a given set $\Omega \in \mathcal{K}_0^n$, the convex set $\tilde{\Omega}$ satisfying that $\Omega = \tilde{\Omega}_t$ and which has the largest possible perimeter. In the case where Ω is a planar polygon this problem has the fairly intuitive solution that $\tilde{\Omega}$ should be the polygon with the same faces as Ω , only moved a distance t along their outward pointing normals. The following lemma tells us that a generalization of this intuitive solution actually works for any possible Ω .

Lemma 2.1. *Let $\Omega \in \mathcal{K}_0^n$ and let Ω^* denote its form body. Then, the maximization problem*

$$\max\{|\partial\tilde{\Omega}| : \tilde{\Omega} \in \mathcal{K}_0^n, \tilde{\Omega}_t = \Omega\}$$

is solved by $\tilde{\Omega} = \Omega + t\Omega^$.*

Proof. Recalling one of the previously mentioned properties of the support function, we have that for any $x \in \partial\Omega = \partial(\tilde{\Omega} \sim tB)$ there exists a $u \in \mathbb{S}^{n-1}$ normal to $\partial\Omega$ at x such that

$$h(\tilde{\Omega} \sim tB, u) = h(\tilde{\Omega}, u) - h(tB, u) = h(\tilde{\Omega}, u) - t.$$

Rearranging this we find for all $u \in \mathcal{U}_0(\Omega)$ that $h(\tilde{\Omega}, u) = h(\Omega, u) + t$. Therefore, it follows that

$$\begin{aligned} \tilde{\Omega} &\subseteq \bigcap_{u \in \mathcal{U}_0(\Omega)} \{x \in \mathbb{R}^n : \langle x, u \rangle \leq h(\Omega, u) + t\} \\ &= \bigcap_{u \in \mathcal{U}_0(\Omega)} \{x \in \mathbb{R}^n : \langle x, u \rangle \leq h(\Omega, u) + h(t\Omega^*, u)\} \\ &= \bigcap_{u \in \mathcal{U}_0(\Omega)} \{x \in \mathbb{R}^n : \langle x, u \rangle \leq h(\Omega + t\Omega^*, u)\} \\ &= \Omega + t\Omega^*. \end{aligned}$$

Thus, since the perimeter is increasing under inclusion of convex bodies, we may conclude that we have the upper bound $|\partial\tilde{\Omega}| \leq |\partial(\Omega + t\Omega^*)|$.

What remains to complete the proof is to show that $\Omega + t\Omega^*$ is an admissible set in the above maximization problem. That $\Omega \subseteq (\Omega + t\Omega^*)_t$ follows from the above argument so we only need to establish the opposite inclusion. Let $x \in \text{reg}(\Omega)$. Then, with u being the unique normal to $\partial\Omega$ at x , we have that

$$h(\Omega + t\Omega^*, u) = h(\Omega, u) + t.$$

Since a convex body can be written as the intersection of its supporting half-spaces we conclude that $x + tu \in (\Omega + t\Omega^*)^c$ implying that $\text{dist}(x, \partial(\Omega + t\Omega^*)) \leq t$. Combining this with the inclusion of Ω in $(\Omega + t\Omega^*)_t$ we find that $\text{reg}(\Omega) \in \partial(\Omega + t\Omega^*)_t$. Since almost every point of $\partial\Omega$ is regular the statement follows. \square

We are now ready to prove our main theorem. By the lemma above we know that any convex body $\tilde{\Omega}$ such that $\tilde{\Omega}_t = \Omega$ satisfies

$$|\partial\tilde{\Omega}| \leq |\partial(\Omega + t\Omega^*)|.$$

Therefore it is sufficient to prove that for any $\Omega \subset \mathbb{R}^n$ with inradius R_{in} it holds that

$$|\partial\Omega| \geq \left(1 - \frac{t}{\tilde{R}_{in}}\right)_+^{n-1} |\partial(\Omega + t\Omega^*)|.$$

Here $\tilde{R}_{in} = R_{in} + t$ is the inner radius of $\Omega + t\Omega^*$. Using the fact that the perimeter of a convex set can be expressed as a mixed volume and the multilinearity of W we find that

$$\begin{aligned} |\partial(\Omega + t\Omega^*)| &= nW(B, \underbrace{\Omega + t\Omega^*, \dots, \Omega + t\Omega^*}_{n-1}) \\ &= n \sum_{m=0}^{n-1} \binom{n-1}{m} t^m W(B, \underbrace{\Omega, \dots, \Omega}_{n-1-m}, \underbrace{\Omega^*, \dots, \Omega^*}_m). \end{aligned}$$

By Lemma 1.5 there exists $x \in \mathbb{R}^n$ such that $x + R_{in}\Omega^* \subseteq \Omega$. Therefore by the translation invariance and the monotonicity of mixed volumes under inclusions we find that

$$\begin{aligned} |\partial(\Omega + t\Omega^*)| &= n \sum_{m=0}^{n-1} \binom{n-1}{m} t^m W(B, \underbrace{\Omega, \dots, \Omega}_{n-1-m}, \underbrace{\Omega^*, \dots, \Omega^*}_m) \\ &= n \sum_{m=0}^{n-1} \binom{n-1}{m} \frac{t^m}{R_{in}^m} W(B, \underbrace{\Omega, \dots, \Omega}_{n-1-m}, \underbrace{R_{in}\Omega^*, \dots, R_{in}\Omega^*}_m) \\ &\leq n \sum_{m=0}^{n-1} \binom{n-1}{m} \frac{t^m}{R_{in}^m} W(B, \underbrace{\Omega, \dots, \Omega}_{n-1}) \\ &= |\partial\Omega| \sum_{m=0}^{n-1} \binom{n-1}{m} \frac{t^m}{R_{in}^m} \\ &= |\partial\Omega| \left(1 + \frac{t}{R_{in}}\right)^{n-1}. \end{aligned}$$

Rearranging and using that $\tilde{R}_{in} = R_{in} + t$ one obtains the desired inequality. Further it is clear from the argument above that equality holds if and only if $\Omega = x + R_{in}\Omega^*$, that is when Ω is homothetic to Ω^* . Thus the proof of Theorem 1.2 is complete.

Deducing Corollary 1.3 is simply a matter of applying the following theorem due to Steinhagen [17]. We note that this theorem appeared in the case of planar convex bodies in earlier work by Blaschke [1], and this simpler case is sufficient for proving the inequality conjectured in [8].

Theorem 2.2 (Steinhagen's inequality). *Let $\Omega \in \mathcal{K}_0^n$ have inradius R_{in} and width ω . Then the following two-sided inequality holds:*

$$\begin{aligned} 2R_{in} \leq \omega &\leq 2\sqrt{n} R_{in} && \text{if } n \text{ is odd,} \\ 2R_{in} \leq \omega &\leq \frac{2(n+1)}{\sqrt{n+2}} R_{in} && \text{if } n \text{ is even.} \end{aligned}$$

The lower bound is attained if Ω is a ball of radius R_{in} . Equality holds in the upper bound if Ω is a regular $(n+1)$ -simplex.

Providing an elementary proof of this theorem, even though it is fairly intuitive, is not an easy task and does not provide much deeper insight into the problem at hand. A complete proof may however be found in [17].

In two dimensions equality in Steinhagen's upper bound holds if and only if Ω is an equilateral triangle. In higher dimensions however one may be able to cut off vertices of the regular simplex without altering the width or inner radius of it. The conjectured uniqueness of the minimizers of the corollary is believed to be true since even though we may slightly modify a regular simplex without altering its width or radius, this would alter the corresponding form body. For equality to hold in the corollary it is necessary that Ω achieves equality in Steinhagen's upper bound and that Ω is homothetic to Ω^* . Therefore, we very much believe that Ω achieving equality in the corollary must be a regular $(n+1)$ -simplex.

Acknowledgements. A great deal of gratitude is owed Professor Ari Laptev for valuable discussions and for proposing the study of the problem at hand. I would also like to thank Professor Bo Berndtsson at Chalmers University of Technology for introducing me to the field of convex geometry.

REFERENCES

- [1] W. Blaschke, *Über den größten Kreis in einer konvexen Punktmenge.*, Jahresbericht der Deutschen Mathematiker-Vereinigung **23** (1914), 369–374.
- [2] G. Bol, *Beweis einer Vermutung von H. Minkowski*, Abh. Math. Sem. Univ. Hamburg **15** (1943), no. 1, 37–56.
- [3] T. Bonnesen and W. Fenchel, *Theorie der konvexen Körper*, Springer, Berlin, 1934. English translation: *Theory of convex bodies*, Edited by L. Boron, C. Christenson and B. Smith, BCS Associates, Moscow, ID, 1987.
- [4] N. S. Brannen, *The Wills conjecture*, Trans. Amer. Math. Soc. **349** (1997), no. 10, 3977–3987.
- [5] A. Colesanti, *A Steiner type formula for convex functions*, Mathematika **44** (1997), no. 1, 195–214.
- [6] A. Dinghas, *Bemerkung zu einer Verschärfung der isoperimetrischen Ungleichung durch H. Hadwiger*, Math. Nachr. **1** (1948), 284–286.
- [7] H. Federer, *Curvature measures*, Trans. Amer. Math. Soc. **93** (1959), 418–491.
- [8] L. Geisinger, A. Laptev, and T. Weidl, *Geometrical Versions of improved Berezin-Li-Yau Inequalities*, Journal of Spectral Theory **1** (2011), 87–109.
- [9] M. A. Hernández Cifre and E. Saorín, *On the volume of inner parallel bodies*, Adv. Geom. **10** (2010), no. 2, 275–286.
- [10] ———, *On inner parallel bodies and quermassintegrals*, Israel J. Math. **177** (2010), 29–47.
- [11] D. Hug, G. Last, and W. Weil, *A local Steiner-type formula for general closed sets and applications*, Math. Z. **246** (2004), no. 1-2, 237–272.
- [12] E. Linke and E. Saorín Gómez, *Decomposition of polytopes using inner parallel bodies*, Monatsh. Math. **176** (2015), no. 4, 575–588.
- [13] G. Matheron, *La formule de Steiner pour les érosions*, J. Appl. Probability **15** (1978), no. 1, 126–135.
- [14] J. R. Sangwine-Yager, *Inner parallel bodies and geometric inequalities*, ProQuest LLC, Ann Arbor, MI, 1978, Thesis (Ph.D.)—University of California, Davis.
- [15] ———, *A Bonnesen-style inradius inequality in 3-space*, Pacific J. Math. **134** (1988), no. 1, 173–178.
- [16] R. Schneider, *Convex bodies: The Brunn-Minkowski theory*, Cambridge Series in Environment and Behavior, Cambridge University Press, 1993.
- [17] P. Steinhagen, *Über die größte Kugel in einer konvexen Punktmenge*, Abh. Math. Sem. Univ. Hamburg **1** (1922), no. 1, 15–26.
- [18] B. Teissier, *Bonnesen-type inequalities in algebraic geometry. I. Introduction to the problem*, Seminar on Differential Geometry, Ann. of Math. Stud., vol. 102, Princeton Univ. Press, Princeton, N.J., 1982, pp. 85–105.
- [19] M. van den Berg, *A uniform bound on trace($e^{t\Delta}$) for convex regions in \mathbb{R}^n with smooth boundaries*, Comm. Math. Phys. **92** (1984), no. 4, 525–530.
- [20] R. Webster, *Convexity*, Oxford science publications, Oxford University Press, 1994.
- [21] H. Weyl, *On the Volume of Tubes*, Amer. J. Math. **61** (1939), no. 2, 461–472.

DEPARTMENT OF MATHEMATICS,
 KTH ROYAL INSTITUTE OF TECHNOLOGY,
 STOCKHOLM, SWEDEN
E-mail address: `simla@kth.se`